

--

--	--	--	--	--

--	--	--	--

0606/11

May/June 2017

2 hours

Additional Materials: Electronic calculator

READ THESE INSTRUCTIONS FIRST

DO **NOT** WRITE IN ANY BARCODES.

You are reminded of the need for clear presentation in your answers.

The total number of marks for this paper is 80.

This document consists of **12** printed pages.

Mathematical Formulae**1. ALGEBRA***Quadratic Equation*

For the equation $ax^2 + bx + c = 0$,

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

Binomial Theorem

$$(a + b)^n = a^n + \binom{n}{1} a^{n-1} b + \binom{n}{2} a^{n-2} b^2 + \dots + \binom{n}{r} a^{n-r} b^r + \dots + b^n,$$

where n is a positive integer and $\binom{n}{r} = \frac{n!}{(n-r)!r!}$

2. TRIGONOMETRY*Identities*

$$\sin^2 A + \cos^2 A = 1$$

$$\sec^2 A = 1 + \tan^2 A$$

$$\operatorname{cosec}^2 A = 1 + \cot^2 A$$

Formulae for ΔABC

$$\frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C}$$

$$a^2 = b^2 + c^2 - 2bc \cos A$$

$$\Delta = \frac{1}{2} bc \sin A$$

1 The line $y = kx - 5$, where k is a positive constant, is a tangent to the curve $y = x^2 + 4x$ at the point A .

(i) Find the exact value of k . [3]

(ii) Find the gradient of the normal to the curve at the point A , giving your answer in the form $a + b\sqrt{5}$, where a and b are constants. [2]

- 2 It is given that $p(x) = x^3 + ax^2 + bx - 48$. When $p(x)$ is divided by $x - 3$ the remainder is 6. Given that $p'(1) = 0$, find the value of a and of b . [5]

- 3 (a) Simplify $\sqrt{x^8 y^{10}} \div \sqrt[3]{x^3 y^{-6}}$, giving your answer in the form $x^a y^b$, where a and b are integers. [2]

- (b) (i) Show that $4(t-2)^{\frac{1}{2}} + 5(t-2)^{\frac{3}{2}}$ can be written in the form $(t-2)^p(qt+r)$, where p , q and r are constants to be found. [3]

- (ii) Hence solve the equation $4(t-2)^{\frac{1}{2}} + 5(t-2)^{\frac{3}{2}} = 0$. [1]


4 (a) It is given that $f(x) = 3e^{-4x} + 5$ for $x \in \mathbb{R}$.

(i) State the range of f . [1]

(ii) Find f^{-1} and state its domain. [4]

(b) It is given that $g(x) = x^2 + 5$ and $h(x) = \ln x$ for $x > 0$. Solve $hg(x) = 2$. [3]

5 (a)



The diagram shows a figure $OACB$, where $\overrightarrow{OA} = \mathbf{a}$, $\overrightarrow{OB} = \mathbf{b}$ and $\overrightarrow{OC} = \mathbf{c}$. The lines AC and OB intersect at the point M where M is the midpoint of the line AC .

(i) Find, in terms of $\mathbf{a}$ and $\mathbf{c}$, the vector $\overrightarrow{OM}$. [2]

(ii) Given that $OM:MB = 2:3$, find $\mathbf{b}$ in terms of $\mathbf{a}$ and $\mathbf{c}$. [2]

(b) Vectors $\mathbf{i}$ and $\mathbf{j}$ are unit vectors parallel to the x -axis and y -axis respectively.


The vector $\mathbf{p}$ has a magnitude of 39 units and has the same direction as $-10\mathbf{i} + 24\mathbf{j}$.

(i) Find $\mathbf{p}$ in terms of $\mathbf{i}$ and $\mathbf{j}$. [2]

(ii) Find the vector $\mathbf{q}$ such that $2\mathbf{p} + \mathbf{q}$ is parallel to the positive y -axis and has a magnitude of 12 units. [3]

(iii) Hence show that $|\mathbf{q}| = k\sqrt{5}$, where k is an integer to be found. [2]

6



The diagram shows a circle, centre O , radius 12 cm . The points A and B lie on the circumference of the circle and form a rectangle with the points C and D . The length of AD is 8 cm and the area of the minor sector AOB is 150 cm^2 .


(i) Show that angle AOB is 2.08 radians, correct to 2 decimal places. [2]

(ii) Find the area of the shaded region $ADCB$. [6]

(iii) Find the perimeter of the shaded region $ADCB$. [3]

- 7 Show that the curve $y = (3x^2 + 8)^{\frac{5}{3}}$ has only one stationary point. Find the coordinates of this stationary point and determine its nature. [8]

- 8 (i) On the axes below sketch the graphs of $y = |2x - 5|$ and $9y = 80x - 16x^2$. [5]



- (ii) Solve $|2x - 5| = 4$. [3]


- (iii) Hence show that the graphs of $y = |2x - 5|$ and $9y = 80x - 16x^2$ intersect at the points where $y = 4$. [1]

- (iv) Hence find the values of x for which $9|2x - 5| \leq 80x - 16x^2$. [2]

9 (i) Show that $5 + 4 \tan^2\left(\frac{x}{3}\right) = 4 \sec^2\left(\frac{x}{3}\right) + 1$. [1]

(ii) Given that $\frac{d}{dx}\left(\tan\left(\frac{x}{3}\right)\right) = \frac{1}{3}\sec^2\left(\frac{x}{3}\right)$, find $\int \sec^2\left(\frac{x}{3}\right) dx$. [1]

(iii)



The diagram shows part of the curve $y = 5 + 4 \tan^2\left(\frac{x}{3}\right)$. Using the results from parts (i) and (ii), find the exact area of the shaded region enclosed by the curve, the x -axis and the lines $x = \frac{\pi}{2}$ and $x = \pi$. [5]

10 (a) Given that $y = \frac{e^{3x}}{4x^2 + 1}$, find $\frac{dy}{dx}$. [3]

(b) Variables x , y and t are such that $y = 4 \cos\left(x + \frac{\pi}{3}\right) + 2\sqrt{3} \sin\left(x + \frac{\pi}{3}\right)$ and $\frac{dy}{dt} = 10$.

(i) Find the value of $\frac{dy}{dx}$ when $x = \frac{\pi}{2}$. [3]

(ii) Find the value of $\frac{dx}{dt}$ when $x = \frac{\pi}{2}$. [2]

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge International Examinations Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cie.org.uk after the live examination series.

Cambridge International Examinations is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.