



CANDIDATE
NAME

--

CENTRE
NUMBER

--	--	--	--	--

CANDIDATE
NUMBER

--	--	--	--

9231/11

May/June 2024

2 hours

You must answer on the question paper.

You will need: List of formulae (MF19)

- Answer **all** questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do **not** use an erasable pen or correction fluid.
- Do **not** write on any bar codes.
- If additional space is needed, you should use the lined page at the end of this booklet; the question number or numbers must be clearly shown.
- You should use a calculator where appropriate.
- You must show all necessary working clearly; no marks will be given for unsupported answers from a calculator.
- Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place for angles in degrees, unless a different level of accuracy is specified in the question.

- The total mark for this paper is 75.
- The number of marks for each question or part question is shown in brackets [].

This document has **16** pages.

- 1 The cubic equation $2x^3 + x^2 - px - 5 = 0$, where p is a positive constant, has roots α, β, γ .

(a) State, in terms of p , the value of $\alpha\beta + \beta\gamma + \gamma\alpha$. [1]

(b) Find the value of $\alpha^2\beta\gamma + \alpha\beta^2\gamma + \alpha\beta\gamma^2$. [2]

This image shows a full page of a handwriting practice worksheet. It consists of multiple sets of three horizontal dotted lines spaced evenly down the page, providing a guide for letter height and placement. The background is plain white, and there are no other markings or text present.

- (d) Given that $\alpha^2 + \beta^2 + \gamma^2 = \frac{1}{3}$, find the value of p . [2]

- 2** Prove by mathematical induction that $6^{4n} + 38^n - 2$ is divisible by 74 for all positive integers n . [6]

[illegible]

- 3 (a)** Use standard results from the list of formulae (MF19) to show that

$$\sum_{r=1}^N r(r+1)(3r+4) = \frac{1}{12}N(N+1)(N+2)(9N+19). \quad [3]$$

This image shows a full page of white paper with horizontal dotted lines. The lines are evenly spaced and run across the width of the page, providing a guide for handwriting or typing. There are no margins, text, or other markings on the page.

- (b)** Express $\frac{3r+4}{r(r+1)}$ in partial fractions and hence use the method of differences to find

$$\sum_{r=1}^N \frac{3r+4}{r(r+1)} \left(\frac{1}{4}\right)^{r+1}$$

in terms of N .

[4]

[illegible]

- (c) Deduce the value of $\sum_{r=1}^{\infty} \frac{3r+4}{r(r+1)} \left(\frac{1}{4}\right)^{r+1}$. [1]

4 The matrix $\mathbf{M}$ is given by $\mathbf{M} = \begin{pmatrix} \frac{1}{2} & -\frac{1}{2}\sqrt{3} \\ \frac{1}{2}\sqrt{3} & \frac{1}{2} \end{pmatrix} \begin{pmatrix} 14 & 0 \\ 0 & 1 \end{pmatrix}$.

(a) The matrix $\mathbf{M}$ represents a sequence of two geometrical transformations in the x - y plane.

Give full details of each transformation, and make clear the order in which they are applied. [4]

[illegible]

(b) Write $\mathbf{M}^{-1}$ as the product of two matrices, neither of which is $\mathbf{I}$. [2]

[illegible]

- (c) Find the equations of the invariant lines, through the origin, of the transformation represented by **M**. [5]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

- (d) The triangle ABC in the x - y plane is transformed by **M** onto triangle DEF .

Given that the area of triangle DEF is 28 cm^2 , find the area of triangle ABC . [2]

.....

.....

.....

.....

.....

.....

5 The points A, B, C have position vectors

$$2\mathbf{i} + 2\mathbf{j} + 4\mathbf{k},$$

$$2\mathbf{i} + 4\mathbf{j} - \mathbf{k},$$

$$-3\mathbf{i} - 3\mathbf{j} + 4\mathbf{k},$$

respectively, relative to the origin O .

(a) Find the equation of the plane ABC , giving your answer in the form $ax + by + cz = d$. [5]

[illegible]

The point D has position vector $2\mathbf{i} + \mathbf{j} + 3\mathbf{k}$.

- (b) Find the perpendicular distance from D to the plane ABC . [2]

- (c) Find the shortest distance between the lines AB and CD . [5]

This image shows a full page of white paper with horizontal dotted lines. The lines are evenly spaced and run across the width of the page, providing a guide for handwriting practice. There are no margins, text, or other markings on the page.

- 6** The curve C has equation $y = \frac{x^2 + ax + 1}{x + 2}$, where $a > \frac{5}{2}$.

(a) Find the equations of the asymptotes of C .

[3]

[illegible]

(b) Show that C has no stationary points.

[4]

This image shows a full page of white paper with horizontal dashed lines, typical of primary-ruled notebook paper. The lines are evenly spaced and run across the width of the page. There are no margins, text, or other markings on the paper.

- (c) Sketch C , stating the coordinates of the point of intersection with the y -axis and labelling the asymptotes. [3]

-
- (d) (i) Sketch the curve with equation $y = \left| \frac{x^2 + ax + 1}{x + 2} \right|$. [2]

- (ii) On your sketch in part (i), draw the line $y = a$. [1]

- (iii) It is given that $\left| \frac{x^2 + ax + 1}{x + 2} \right| < a$ for $-5 - \sqrt{14} < x < -3$ and $-5 + \sqrt{14} < x < 3$.

Find the value of a . [2]

.....

.....

.....

.....

.....

.....

.....

7 The curve C has polar equation $r^2 = (\pi - \theta) \tan^{-1}(\pi - \theta)$, for $0 \leq \theta \leq \pi$.

- (a) Sketch C and state the polar coordinates of the point of C furthest from the pole. [3]

.....

- (b) Using the substitution $u = \pi - \theta$, or otherwise, find the area of the region enclosed by C and the initial line. [7]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

- (c)** Show that, at the point of C furthest from the initial line,

$$2(\pi - \theta) \tan^{-1}(\pi - \theta) \cot \theta - \frac{\pi - \theta}{1 + (\pi - \theta)^2} - \tan^{-1}(\pi - \theta) = 0$$

and verify that this equation has a root for θ between 1.2 and 1.3.

[5]

[illegible]

[illegible]

Cambridge Assessment International Education is part of Cambridge Assessment. Cambridge Assessment is the brand name of the University of Cambridge Local Examinations Syndicate (UCLES), which is a department of the University of Cambridge.