

June 2004

GCE AS LEVEL

MARK SCHEME

MAXIMUM MARK: 50

SYLLABUS/COMPONENT: 9709/02

**MATHEMATICS
Paper 2 (Pure 2)**



Page 1	Mark Scheme	Syllabus	Paper
	A AND AS LEVEL – JUNE 2004	9709	2

1	Use logarithms to linearise an equation	M1	
	Obtain $\frac{x}{y} = \frac{\ln 5}{\ln 2}$ or equivalent	A1	
	Obtain answer 2.32	A1	3
2	(i) Use the given iterative formula correctly at least ONCE with $x_1 = 3$	M1	
	Obtain final answer 3.142	A1	
	Show sufficient iterations to justify its accuracy to 3 d.p.	A1	3
	(ii) State any suitable equation e.g. $x = \frac{1}{5} \left(4x + \frac{306}{x^4} \right)$	B1	
	Derive the given answer α (or x) = $\sqrt[5]{306}$	B1	2
3	(i) Substitute $x = 3$ and equate to zero	M1	
	Obtain answer $\alpha = -1$	A1	2
	(ii) At any stage, state that $x = 3$ is a solution	B1	
	EITHER: Attempt division by $(x-3)$ reaching a partial quotient of $2x^2 + kx$	M1	
	Obtain quadratic factor $2x^2 + 5x + 2$	A1	
	Obtain solutions $x = -2$ and $x = -\frac{1}{2}$	A1	
	OR: Obtain solution $x = -2$ by trial and error	B1	
	Obtain solution $x = -\frac{1}{2}$ similarly	B2	4
	[If an attempt at the quadratic factor is made by inspection, the M1 is earned if it reaches an unknown factor of $2x^2 + bx + c$ and an equation in b and/or c .]		
4	(i) State answer $R = 5$	B1	
	Use trigonometric formulae to find α	M1	
	Obtain answer $\alpha = 53.13^\circ$	A1	3
	(ii) Carry out, or indicate need for, calculation of $\sin^{-1}(4.5/5)$	M1	
	Obtain answer 11.0°	A1√	
	Carry out correct method for the second root e.g. $180^\circ - 64.16^\circ - 53.13^\circ$	M1	
	Obtain answer 62.7° and no others in the range	A1√	4
	[Ignore answers outside the given range.]		
	(iii) State least value is 2	B1√	1
5	(i) State derivative of the form $(e^{-x} \pm xe^{-x})$. Allow $xe^x \pm e^x$ {via quotient rule}	M1	
	Obtain correct derivative of $e^{\pm x} - xe^{-x}$	A1	
	Equate derivative to zero and solve for x	M1	
	Obtain answer $x = 1$	A1	4
	(ii) Show or imply correct ordinates 0, 0.367879..., 0.27067...	B1	
	Use correct formula, or equivalent, with $h = 1$ and three ordinates	M1	
	Obtain answer 0.50 with no errors seen	A1	3
	(iii) Justify statement that the rule gives an under-estimate	B1	1

Page 2	Mark Scheme	Syllabus	Paper
	A AND AS LEVEL – JUNE 2004	9709	2

- 6 (i) State that $\frac{dx}{dt} = 2 + \frac{1}{t}$ or $\frac{dy}{dt} = 1 - \frac{4}{t^2}$, or equivalent B1
- Use $\frac{dy}{dx} = \frac{dy}{dt} \div \frac{dx}{dt}$ M1
- Obtain the given answer A1 **3**
- (ii) Substitute $t = 1$ in $\frac{dy}{dx}$ and both parametric equations M1
- Obtain $\frac{dy}{dx} = -1$ and coordinates (2, 5) A1
- State equation of tangent in any correct horizontal form e.g. $x + y = 7$ A1√ **3**
- (iii) Equate $\frac{dy}{dx}$ to zero and solve for t M1
- Obtain answer $t = 2$ A1
- Obtain answer $y = 4$ A1
- Show by any method (but not via $\frac{d}{dt}(y')$) that this is a minimum point A1 **4**
- 7 (i) Make relevant use of the $\cos(A + B)$ formula M1*
- Make relevant use of $\cos 2A$ and $\sin 2A$ formulae M1*
- Obtain a correct expression in terms of $\cos A$ and $\sin A$ A1
- Use $\sin^2 A = 1 - \cos^2 A$ to obtain an expression in terms of $\cos A$ M1(dep*)
- Obtain given answer correctly A1 **5**
- (ii) Replace integrand by $\frac{1}{4} \cos 3x + \frac{3}{4} \cos x$, or equivalent B1
- Integrate, obtaining $\frac{1}{12} \sin 3x + \frac{3}{4} \sin x$, or equivalent B1 + B1√
- Use limits correctly M1
- Obtain given answer A1 **5**